

**PRE-OPERATIVE MEDICATIONS POLICIES**
**ADHD DRUGS – Hold Day of Surgery**

Common ADHD Drugs	<ul style="list-style-type: none"> <li>• <b>Adderall</b> (Amphetamine/Dextroamphetamine)</li> <li>• <b>Concerta</b> (Methylphenidate)</li> <li>• <b>Daytrana</b> (Methylphenidate)</li> <li>• <b>Desoxyn</b> (Methamphetamine)</li> <li>• <b>Dexedrine</b> (Dextroamphetamine)</li> <li>• <b>Dextrostat</b> (Dextroamphetamine)</li> <li>• <b>Focalin</b> (Dexmethylphenidate)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Intuniv</b> (Guanfacine)</li> <li>• <b>Metadate</b> (Methylphenidate)</li> <li>• <b>Methylin</b> (Methylphenidate)</li> <li>• <b>Ritalin</b> (Methylphenidate)</li> <li>• <b>Strattera</b> (Atomoxetine)</li> <li>• <b>Tenex</b> (Guanfacine)</li> <li>• <b>Vyvanse</b> (Lisdexamfetamine)</li> </ul>
-------------------	---	---

**WEIGHT-LOSS DRUGS**
**Appetite Suppressants – Hold for a Week**

Common Appetite Suppressants	<ul style="list-style-type: none"> <li>• <b>Phentermine</b></li> <li>• <b>Phendimetrazine</b></li> </ul>
------------------------------	--

**GL-1 Receptor Agonists – Hold Daily Medications Day of Surgery | Hold Weekly Medications for a Week**

Common GLP-1 Agonists	<ul style="list-style-type: none"> <li>• <b>Ozempic</b> (Semaglutide)</li> <li>• <b>Rybelsus</b> (Semaglutide)</li> <li>• <b>Trulicity</b> (Dulaglutide)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Victoza</b> (Liraglutide)</li> <li>• <b>Wegovy</b> (Semaglutide)</li> </ul>
-----------------------	---	---

**ANTICOAGULANT/ANTIPLATELET DRUGS – Hold as Instructed by Prescribing Physician**

Common Anticoagulants & Antiplatelets	<ul style="list-style-type: none"> <li>• <b>Aggrenox</b> (Acetylsalicylic Acid/Dipyridamole)</li> <li>• <b>Arixtra</b> (Fondaparinux)</li> <li>• <b>Brilinta</b> (Ticagrelor)</li> <li>• <b>Coumadin</b> (Warfarin)</li> <li>• <b>Effient</b> (Prasugrel)</li> <li>• <b>Eliquis</b> (Apixaban)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Jantoven</b> (Warfarin)</li> <li>• <b>Plavix</b> (Clopidogrel)</li> <li>• <b>Pletal</b> (Cilostazol)</li> <li>• <b>Pradaxa</b> (Dabigatran)</li> <li>• <b>Ticlid</b> (Ticlopidine)</li> <li>• <b>Xarelto</b> (Rivaroxaban)</li> </ul>
---------------------------------------	---	---

**ACE INHIBITORS – Hold Day of Surgery**

Common Ace Inhibitors	<ul style="list-style-type: none"> <li>• <b>Accupril</b> (Quinapril)</li> <li>• <b>Aceon</b> (Perindopril)</li> <li>• <b>Altace</b> (Ramipril)</li> <li>• <b>Capoten</b> (Captopril)</li> <li>• <b>Lotensin</b> (Benazepril)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Monopril</b> (Fosinopril)</li> <li>• <b>Prinivil</b> (Lisinopril)</li> <li>• <b>Univasc</b> (Moexipril)</li> <li>• <b>Vasotec</b> (Enalapril)</li> <li>• <b>Zestril</b> (Lisinopril)</li> </ul>
-----------------------	---	---

**PHOSPHODIESTERASE-5 INHIBITORS – Hold for 36 Hours**

Common Phosphodiesterase-5 Inhibitors	<ul style="list-style-type: none"> <li>• <b>Cialis</b> (Tadalafil)</li> <li>• <b>Levitra</b> (Vardenafil)</li> <li>• <b>Mvix</b> (Mirodenafil)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Viagra</b> (Sildenafil)</li> <li>• <b>Zydena</b> (Udenafil)</li> </ul>
---------------------------------------	---	--

**DIABETES MEDICATIONS**

- **Oral Hypoglycemics – Hold Day of Surgery**
- **Rapid-Acting or Short-Acting Insulin** (e.g., regular, lispro, aspart, glulisine) – **Hold Day of Surgery**
- **Intermediate-Acting Insulin** (e.g., NPH) – **Take Half AM Dose**
- **Long-Acting Insulin** (e.g., glargine, detemir) – **Take Half AM Dose**
- **Pump – Continue at Same Rate for Procedures < 2 Hours | Convert to Infusion for Longer Procedures**

**NALTREXONE – Hold One Week | Hold 4 Weeks if IM Injection**